Fisheries and Oceans Canada | Pêches et Océans Canada

Canadian Stock Assessment Secretariat
Research Document 99/200

Not to be cited without
permission of the authors[1]

Secrétariat canadien pour l'évaluation des stocks
Document de recherche 99/200

Ne pas citer sans
autorisation des auteurs[1]

# Pacific Hake - Strait of Georgia Stock Assessment for 1999 and Recommended Yield Options for 2000

M.W. Saunders and G.A. McFarlane

Department of Fisheries and Oceans
Biological Sciences Branch
Pacific Biological Station
3190 Hammond Bay Road
Nanaimo, B.C. V9R 5K6

[1] This series documents the scientific basis for the evaluation of fisheries resources in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research documents are produced in the official language in which they are provided to the Secretariat.

[1] La présente série documente les bases scientifiques des évaluations des ressources halieutiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au secrétariat.

ISSN 1480-4883
Ottawa, 1999
Canada

## ABSTRACT

The fishery in the Strait of Georgia decreased substantially to 3,996t in 1998 from 6,561t in 1997, a result of low market demand. Age and growth data continue to indicate strong recruitment during the 1990's and a coincidental decline in the mean size-at-age. Yield options remain unchanged from the previous assessment with a range of 7,554-14,687 t. Due to uncertainty in the current assessment, including evidence of increasing seal predation we recommend that managers choose a quota from the lower half of the yield range.

## RÉSUMÉ

La pêcherie dans le détroit de Géorgie a diminué de façon appréciable pour passer de 6 561 t en 1997 à 3 996 t en 1998 suite à une réduction de la demande. Les données sur l'âge et la croissance continuent d'indiquer un fort recrutement au cours des années 1990 et un déclin simultané de la taille moyenne selon l'âge. La gamme des options de rendement, qui se situe entre 7 554 t et 14 687 t, demeure inchangée par rapport à l'évaluation précédente. Étant donné le caractère incertain de la présente évaluation, notamment des indices d'une prédation accrue par les phoques, nous recommandons que les gestionnaires fixent le quota à une valeur se situant dans la demie inférieure de la gamme de rendement.

## 1.0 Introduction

This is an interim assessment for Strait of Georgia Pacific hake and yield options are not revised from the previous assessment (Saunders and McFarlane 1999). We review current fishery and biological data.

## 1.1 Landing Statistics

Pacific hake is the most abundant resident fish in the Strait of Georgia. Landings of hake from the Strait of Georgia have gradually increased from the inception of the fishery in 1978. The quota was attained for the first time in 1995 when 11,859 t were landed (Table 1.1). Since then the catch has declined substantially to 3,996t in 1998, despite higher available quotas (Table 1.1). Fishers indicated that the lower catches reflect depressed market demand at the time of peak abundance, reduced availability of hake in the post-spawning period and less interest due to an increasingly poor recovery rate due to continuously declining size of fish since 1990.

## 1.2. Condition of the Stock

### Biological information

Age composition data (Fig. 1.1) indicate that pulses of recruitment have occurred every 3-5 years since monitoring began. The dominant year-classes are 77-78, 83-84,87, 91-92, 95, and 98-99. Since 1994 there has been a dramatic decline in the mean size at age (Fig. 1.2) for adult hake. For example, 4 year old females (age at full recruitment) throughout the 1980's were approximately 40 cm in length and have gradually declined to 35 cm by 1995 (Beamish et al. In press). The size at age of age 1's and 2's have remained constant, if not slightly increasing over the time series (Fig. 1.2). The most heavily affected year of growth appears to be the third year, a period when hake are becoming mature and moving deeper in the water column to feed. The amount of growth occurring during years 4 through 6 appears constant over the time series (Fig 1.2) indicating that cohorts do not compensate in later years for the reduced growth at age 3. This is illustrated further by growth trajectories for dominant year-classes presented in Fig. 1.3. The 1991 and 1992 cohorts are 5cm smaller after age 3 than the fish from earlier years. Finally it should be noted that there appears to be a slight increase for all but age 2's in 1999 (Fig 1.2) .

It is clear that the dynamics of the Pacific hake stock in the St. of Georgia have changed since the early 1990's. At present the mechanism for these changes is unknown however it is coincident with changes in the oceanography of the Strait, particularly water temperature, timing of the spring discharge of the Fraser River, and timing of the spring plankton bloom (Beamish & McFarlane In press). The relationship between climate, ocean processes and productivity of the Strait are currently being investigated.

Another factor to consider is predation by harbour seal populations. A recent examination of harbour seal populations in the Strait of Georgia found that the population had increased

from 4,000 seals in 1973 to about 41,000 (95% confidence interval 30,000-52,500) in 1996 (Olesiuk 1999). The annual growth rate of the population was about 11.9% through the 1970's and 80's and slowed to 7.2% in the early to mid 1990's (Olesiuk 1999). There are indications that the population has been stable since 1996 (Olesiuk 1999). Rates of hake consumption were estimated from scat analysis conducted from 1982-88 (Olesiuk 1993). Hake represented 42.6 percent of the diet by weight and annual consumption was estimated to be between 2,215 and 6,664t. At the 1996 population level, this population is estimated to consume approximately 11,000t of hake annually although given the width of confidence intervals for abundance and consumption, the actual amount could vary between 4,400 and 21,000t. A large amount of the uncertainty lies with the assumption that the rates of consumption developed in the 1980's apply to the current population. In addition, the population of seals in 1980's contained a large number of juveniles and was distributed quite differently from the present. Continued monitoring of abundance, more current assessment of the feeding habits of seals in the Strait of Georgia and examination of the size and age component component of hake stock impacted would be helpful in assessing the impact of seals on hake.

Hydroacoustic surveys

The development of an age structured assessment hinges on development of an appropriate index of abundance. Hydroacoustic and trawl swept-volume survey techniques were applied to hake in 1981 and produced similar results with biomass estimated to range between 85,000 and 150,000t (McFarlane and Beamish, 1985). Since then hydroacoustic surveys of the spawning population have been conducted in the springs of 1998, 1993, and annually from 1996 through 1998. During that period a number of changes to the hydroacoustic method, including the development of length-based methods of applying target strength and implementation of new hardware, made interpretation of the time series difficult. The time series was standardized in 1999 by Kieser et al. (1999) and is presented in Table 1.2.

While the application of these hydroacoustic techniques was deemed sound there were are a number of aspects related to survey timing that indicate surveys conducted during April are overestimates of biomass and that recent surveys conducted during February/March are underestimates. The three surveys conducted during the spring of 1981 indicated that biomass available to the survey increased from January to April when peak spawning occurred (Table 1.2). Unfortunately during the April survey, signal from the spring plankton bloom was confounded with the hake producing an overestimate of hake. This occurred again in 1993 when a survey was conducted in mid-March. In order to avoid the plankton surveys conducted since 1993 were carried out in February. Unfortunately, based on the state of maturity and sex ratio, the surveys conducted from 1996-98 appear to have been well in advance of the peak in spawning (Kieser et al. 1999) when biomass would be expected to be higher.

At the present time we do not have an adequate absolute or relative index, of stock size and the recent biomass estimates should be considered a conservative minimum. Given the conservative bias resulting from factors discussed above and reported in detail in Keiser et al. (1999) we believe the biomass of hake in the 1990's to be stable at approximately 50-60,000t. In the future, as

recommended by Kieser et al. (1999) a number of steps should be taken to improve the survey. Since the new split beam/multifrequency EK500 sounder used since 1996 has the potential to separate the plankton signal from the hake the feasibility of April or summer surveys when biomass availability is high should be examined. In addition multiple surveys should be conducted within one year to capture peak biomass and surveys should examine inlets adjacent to the Strait of Georgia to confirm that the majority of adult hake are in the survey area.

In the previous assessment (Saunders and McFarlane 1999) low-high risk annual rates of fishing mortality generated for the offshore (west coast of Vancouver Island) stock (Dorn 1996) were applied to the 1997 estimate of exploitable biomass resulting in a range in yield from 7,554- 14,687 t. Re-calculating using the 1998 survey estimate of 33,681t would produce a recommended range of 6,063t-11,788t. There was however, evidence based on a sex ratio that favoured females and a high proportion of pre-spawning fish (Kieser et. al 1999) that this survey, like the 1996 and 1997 surveys, is biased conservatively due to survey timing. The yield range presented in the previous assessment is close to the median of the three most recent surveys that are all believed to be underestimates. We recommend leaving the yield range unchanged from the previous assessment.

## 1.3. Yield Options

The apparent stability in biomass in spite of decreased growth implies increased recruitment in recent years. Had growth remained comparable to levels experienced in the 1980's, the numbers present in the 1990's would likely equate to close to 100Kt. The range in yield options are unchanged from the previous assessment 7,554-14,687t. We recommend that until a full assessment is completed and factors such as decreasing size-at-age and marine mammal predation are explicitly considered, that managers choose from the lower half of the yield range.

## 1.4. Literature Cited

Beamish, R.J. and G.A. McFarlane. In press. Applying ecosystem management to fisheries in the Strait of Georgia. *In* Ecosystem considerations in Fisheries Management. Lowell Wakefield Symposium Series.

Beamish, R.J. and G.A. McFarlane and J.R. King. In press. Fisheries climatology: understanding decadal scale processes that naturally regulate British Columbia fish populations. *In* Fisheries Oceanography: a science for the new millennium. T. Parsons and P. Harrison (eds.).

Dorn, M. 1996. Status of the coastal Pacific whiting resource in 1996. *In* Pacific Fishery Management Council, Appendix Volume 1: Status of the Pacific Coast groundfish fishery through 1996 and recommended acceptable biological catches in 1997.

Gulland, J.A. 1970. The fish resources of the ocean. FAO Fish. Tech. Paper No. 97: 425 p.

Kieser, R., M.W. Saunders and K. Cooke. 1999. Review of hydroacoustic methodology and Pacific hake biomass estimates for the Strait of Georgia, 1981 to 1998. Canadian Stock Assessment Secretariat Research Document 99/15.

Olesiuk, P.F. 1993. Annual prey consumption by harbour seals (*Phoca vitulina*) in the Strait of Georgia, British Columbia. 91: 491-515.

Olesiuk, P.F. 1999. Status of harbour seals (*Phoca vitulina*) in British Columbia. Canadian Stock Assessment Secretariat Research Document 99/33.

McFarlane, G.A. and R.J. Beamish. 1985. Biology and fishery of Pacific whiting *Merluccius productus*, in the Strait of Georgia. Marine Fisheries Review 47(2).

Saunders, M.W. and G.A. McFarlane. 1999. Pacific hake – Strait of Georgia stock assessment for 1998 and recommended yield options for 1999. PSARC Working Paper G98-2

Table 1.1. History of Strait of Georgia Pacific hake scientific advice to managers, management quota and landings (t), excluding dumped and discarded fish from 1978-1999. Sources: 1978-95 DFO Science Branch catch and effort database (Pacific Biological Station, Nanaimo, B.C.); 1996-1999 Archipelago Marine Research annual Dock Monitoring Program summary.

| Year | TAC (t) | Basis for advice | Quota (t) | Landings (t) |
|---|---|---|---|---|
| 1976 | - | - | - | 5 |
| 1977 | - | - | - | 0 |
| 1978 | - | - | - | 2 |
| 1979 | - | - | - | 516 |
| 1980 | 10,000 | Gulland (1970) MSY using swept-volume and acoustic survey estimates of $B_0$. | - | 508 |
| 1981 | 10,000 | " | 10,000 | 2236 |
| 1982 | 10,000 | " | 10,000 | 2786 |
| 1983 | 10,000 | " | 10,000 | 3122 |
| 1984 | 10,000 | " | 10,000 | 4391 |
| 1985 | 10-15,000 | As above but range considered the plausible range in B and M. | 10,000 | 4886 |
| 1986 | 10-15,000 | " | 10,000 | 4798 |
| 1987 | 10-15,000 | " | 10,000 | 8161 |
| 1988 | 8-24,000 | VPA conducted with constant yield (8-16,000t) and variable yield (16-24,000t projections. | 11,000 | 5254 |
| 1989 | 8-14,000 | VPA conducted with constant yield (8-16,000t projections. | 11,000 | 6299 |
| 1990 | 8-14,000 | " | 11,000 | 6810 |
| 1991 | 8-14,000 | " | 11,000 | 7324 |
| 1992 | 8-14,000 | " | 11,000 | 8485 |
| 1993 | 8-14,000 | " | 11,000 | 4368 |
| 1994 | 8-14,000 | " | 11,000 | 9695 |
| 1995 | 8-14,000 | " | 11,000 | 11859 |
| 1996 | 8-14,000 | " | 11,000 | 6583 |
| 1997 | 10,300-20,100 | Low risk $F_{opt}$ from offshore stock applied to the 1996 biomass estimate. | 15,200 | 6561 |
| 1998 | 7,500-14,700 | Low risk $F_{opt}$ from offshore stock applied to the 1997 biomass estimate. | 14,687 | 3996 |
| 1999 | 7,500-14,700 | " | 10,000 | |

Table 1.2 Summary of biomass estimates of Strait of Georgia Pacific hake from hydroacoustic surveys conducted from 1981 to 1998 (Kieser et al. 1999).

| Date | | 1981 Jan 12-23 | 1981 Feb 09-20 | 1981 April 13-24 | 1981 April 13-24 | 1988 March 18-28 | 1993 March 08-25 | 1996 Feb 20-05 | 1997 Feb 17-28 | 1998 Feb 16-26 |
|---|---|---|---|---|---|---|---|---|---|---|
| Vessel | | GBR | GBR | GBR | GBR | WER | WER | WER | WER | WER |
| Sounder | | BioSonics | BioSonics | BioSonics | BioSonics | BioSonics | BioSonics | EK500 | EK500 | EK500 |
| Comment | | | | Measured | Adjusted | | Fishing March 18-25 | | | |
| | Area | Biomass, t | | | | | | | | |
| Shallow | South | 2060 | 3401 | 36010 | 4573 | 10759 | 5176 | 2294 | 1102 | 7 |
| | North | 4532 | 4258 | 10268 | 1769 | 0 | 9655 | 1660 | 1687 | 0 |
| | NA | 2054 | 2129 | 2311 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Malaspina | 7004 | 3491 | 10715 | 0 | 0 | 438 | 609 | 346 | 0 |
| | Sabine | 198 | 1325 | 1737 | 0 | 0 | 195 | 0 | 63 | 0 |
| | **Total** | **15848** | **14604** | **61041** | **6343** | **10759** | **15465** | **4564** | **3198** | **7** |
| Deep | South | 20904 | 38239 | 49104 | 49104 | 55415 | 55773 | 34556 | 26731 | 25120 |
| | North | 10715 | 15572 | 5110 | 5110 | 0 | 23765 | 19073 | 13071 | 4242 |
| | NA | 946 | 755 | 709 | 709 | 0 | 0 | 0 | 0 | 0 |
| | Malaspina | 4095 | 8733 | 7075 | 7075 | 0 | 6416 | 2072 | 3169 | 2660 |
| | Sabine | 879 | 2622 | 3201 | 3201 | 0 | 3590 | 0 | 355 | 1653 |
| | **Total** | **37539** | **65921** | **65199** | **65199** | **55415** | **89543** | **55702** | **43326** | **33674** |
| Shallow and Deep | South | 22964 | 41640 | 85114 | 53677 | 66174 | 60949 | 36851 | 27833 | 25127 |
| | North | 15247 | 19830 | 15378 | 6880 | 0 | 33420 | 20734 | 14758 | 4242 |
| | NA | 3000 | 2884 | 3019 | 709 | 0 | 0 | 0 | 0 | 0 |
| | Malaspina | 11099 | 12224 | 17791 | 7075 | 0 | 6855 | 2682 | 3515 | 2660 |
| | Sabine | 1076 | 3947 | 4938 | 3201 | 0 | 3786 | 0 | 418 | 1653 |
| | **Total** | **53387** | **80525** | **126240** | **71542** | **66174** | **105008** | **60266** | **46524** | **33681** |




Figure 1.1. Age compositions from Strait of Georgia Pacific hake from 1976-98.




Figure 1.2. Mean size at age of Strait of Georgia Pacific hake from 1976-99.




Figure 1.3. Mean size at age of prominent cohorts of Strait of Georgia Pacific hake from 1978-92.